

***REVERSIBLE WETTING OF NaCl NANOPARTICLES AT RELATIVE HUMIDITIES BELOW DELIQUESCENCE OBSERVED BY ENVIRONMENTAL NON-CONTACT AFM***

D. A. Bruzewicz<sup>2</sup>, A. Checco<sup>2</sup>, B. M. Ocko<sup>2</sup>, E. Lewis<sup>1</sup>, R. L. McGraw<sup>1</sup>, and S. E. Schwartz<sup>1</sup>

<sup>1</sup> Atmospheric Sciences Division, Brookhaven National Laboratory, Upton, NY 11973-5000.

<sup>2</sup> Condensed Matter Physics and Materials Science Department, Brookhaven National Laboratory, Upton, NY 11973-5000.

*For presentation at the*  
2009 American Geophysical Union Fall Meeting  
San Francisco, CA  
Dec. 14-18, 2009

**Environmental Sciences Department/Atmospheric Sciences Division**  
**Brookhaven National Laboratory**  
P.O. Box, Upton, NY  
[www.bnl.gov](http://www.bnl.gov)

**ABSTRACT**

The behavior of NaCl nanoparticles as a function of relative humidity (RH) was characterized by depositing particles on a prepared hydrophobic surface and measuring their height via non-contact environmental atomic force microscopy (AFM). Non-contact AFM allows greater sensitivity to changes in the size of particles than does contact AFM or scanning electron microscopy, and greater sensitivity to changes in shape than do mass-based techniques. Crystalline cubic NaCl nanoparticles with sides of 35 to 150 nm were found to reversibly take up water with increasing RH, and to form a liquid-like surface layer of thickness 2 to 4 nm at humidities well below the deliquescence point of 75.0% at 20°C. Measurable uptake begins at 70% RH. The maximum thickness of the layer increases with increasing RH for a given particle size and, for a given RH, increases with increasing particle size over the range studied. The liquid-like behavior of the layer is indicated by a reversible “rounding” at the tops of the particles, where the ratio of particle height to radius of curvature increases from zero (flat top) at 68% RH to 0.7 at 74% RH. These observations suggest that a reorganization of mass occurs on the solid NaCl nanoparticle, and hence that the behavior of NaCl aerosol nanoparticles at RH between 70 and 75% RH is more complex than an abrupt first-order phase transition. Theoretical treatments of the phase transition should therefore account for both the presence of a liquid-like layer prior to deliquescence, and the RH-dependent thickness of the layer.

---

**NOTICE:** This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.